

*Current Concepts***CARBON MONOXIDE POISONING**

ARMIN ERNST, M.D., AND JOSEPH D. ZIBRAK, M.D.

CARBON monoxide intoxication continues to be one of the most common causes of morbidity due to poisoning in the United States.^{1,2} It may be intentional or accidental, and exposure may be lethal. Approximately 600 accidental deaths due to carbon monoxide poisoning are reported annually in the United States,³ and the number of intentional carbon monoxide-related deaths is 5 to 10 times higher.¹ The rate of accidental death caused by carbon monoxide from motor vehicles is higher in the northern United States and peaks during the winter months.⁴ The intentional deaths occur year-round without significant peaks.¹ The severe winter of 1995–1996 was associated with increased numbers of reported injuries from carbon monoxide exposure.^{5,6} In the winter of 1997–1998, the unusually high number of deaths from carbon monoxide was related to the use of poorly ventilated gasoline-powered generators during a severe ice storm in the northeastern United States.

SOURCES OF CARBON MONOXIDE

Carbon monoxide is a product of the incomplete combustion of hydrocarbons. The concentration of carbon monoxide in the atmosphere is usually less than 0.001 percent. The levels are higher in urban areas than in rural areas. Endogenous carbon monoxide production from the catabolism of hemoglobin is a component of normal biochemical processes. A low base-line level of carboxyhemoglobin is detectable in every person. Tobacco smoke is an important source of carbon monoxide. Blood carboxyhemoglobin commonly reaches a level of 10 percent in smokers and may even exceed 15 percent, as compared with 1 to 3 percent in nonsmokers.⁷⁻⁹

From the Division of Pulmonary and Critical Care Medicine, Department of Medicine, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston. Address reprint requests to Dr. Ernst at the Division of Pulmonary and Critical Care Medicine, Palmer Bldg., Rm. 108, Beth Israel Deaconess Medical Center, West Campus, 1 Deaconess Rd., Boston, MA 02215.

©1998, Massachusetts Medical Society.

The sources of exogenous carbon monoxide that cause poisoning include motor vehicle exhaust fumes, poorly functioning heating systems, and inhaled smoke.¹⁰ Propane-operated forklifts have been implicated as a cause of headache in warehouse workers.¹¹ “Cleaner” fuels such as propane and methane undergo more complete combustion but have also been reported to be sources of carbon monoxide poisoning.

The carbon monoxide in motor vehicle exhaust fumes accounts for the majority of deaths from carbon monoxide poisoning in the United States.¹² Of the 11,547 accidental carbon monoxide deaths reported between 1979 and 1988, motor vehicle exhaust accounted for 57 percent.¹ In a series of 56 motor vehicle-associated deaths reported from 1980 to 1995, 43 percent were due to faulty exhaust systems, 39 percent to operation in an improperly ventilated structure, and 18 percent to the use of a fuel-burning heating device in the passenger compartment.⁴ Lethal concentrations of carboxyhemoglobin can be achieved within 10 minutes in the confines of a closed garage.¹³ Carbon monoxide from motor vehicles can also cause death in semienlosed spaces or in working or living quarters adjacent to garages.¹²

An often overlooked source of carbon monoxide poisoning is methylene chloride, a common component of paint remover and other solvents. Methylene chloride is readily absorbed through the skin and lungs as a vapor and circulates to the liver, where its metabolism results in the generation of carbon monoxide.¹⁴

PATHOPHYSIOLOGY

Carbon monoxide is a colorless, odorless, and nonirritant toxic gas that is easily absorbed through the lungs. The amount of gas absorbed is dependent on the minute ventilation, the duration of exposure, and the relative concentrations of carbon monoxide and oxygen in the environment.¹⁵ Carbon monoxide is principally eliminated by the lungs as an unchanged gas. Less than 1 percent is oxidized to carbon dioxide.¹⁶ Ten to 15 percent of carbon monoxide is bound to proteins, including myoglobin and cytochrome-*c* oxidase.¹⁷ Less than 1 percent of the absorbed gas exists in solution.

Carbon monoxide toxicity appears to result from a combination of tissue hypoxia and direct carbon monoxide-mediated damage at the cellular level. Carbon monoxide competes with oxygen for binding to hemoglobin. The affinity of hemoglobin for carbon monoxide is 200 to 250 times as great as its affinity for oxygen.¹⁸ The consequences of this competitive binding are a shift of the oxygen-hemoglobin

bin dissociation curve to the left and its alteration to a more hyperbolic shape (Fig. 1). These alterations result in impaired release of oxygen at the tissue level and cellular hypoxia.¹⁹ The binding of carbon monoxide to hemoglobin alone does not account for all of the pathophysiologic consequences observed. In studies in animals, transfusion of blood with highly saturated carboxyhemoglobin but minimal free carbon monoxide does not reproducibly result in clinical symptoms.²⁰ This observation suggests that the small fraction of free carbon monoxide dissolved in plasma has an important role.

Recent investigations suggest other mechanisms of carbon monoxide-mediated toxicity. One hypothesis is that carbon monoxide-induced tissue hypoxia may be followed by reoxygenation injury to the central nervous system. Hyperoxygenation facilitates the production of partially reduced oxygen species, which in turn can oxidize essential proteins and nucleic acids, resulting in typical reperfusion injury.²¹ In addition, carbon monoxide exposure has been shown to cause lipid peroxidation (degradation of unsaturated fatty acids), leading to reversible demyelination of




Figure 1. Oxygen-Hemoglobin Dissociation Curve.

The presence of carboxyhemoglobin shifts the curve to the left and changes it to a more hyperbolic shape. This results in a decrease in oxygen-carrying capacity and impaired release of oxygen at the tissue level.

central nervous system lipids.²² Carbon monoxide exposure also creates substantial oxidative stress on cells, with production of oxygen radicals resulting from the conversion of xanthine dehydrogenase to xanthine oxidase.²³

Carbon monoxide exposure has an especially deleterious effect on pregnant women, because of the greater sensitivity of the fetus to the harmful effects of the gas. Data from studies in animals suggest a significant lag time in carbon monoxide uptake between mother and fetus. Fetal steady states can occur up to 40 hours after maternal steady states are achieved. The final carboxyhemoglobin levels in the fetus may significantly exceed the levels in the mother.²⁴ The exaggerated leftward shift of fetal carboxyhemoglobin makes tissue hypoxia more severe by causing less oxygen to be released to fetal tissues.⁷ Although the teratogenicity of carbon monoxide is controversial, the risk of fetal injury seems to be increased by carbon monoxide.²⁵⁻²⁷

CLINICAL SIGNS AND SYMPTOMS

The clinical symptoms of carbon monoxide poisoning are nonspecific and can suggest a broad range of diagnostic possibilities. The signs and symptoms of nonlethal carbon monoxide exposure may mimic those of a nonspecific viral illness. Since viral illnesses and carbon monoxide exposure both peak during the winter, a substantial number of initial misdiagnoses may occur.²⁸ Carbon monoxide poisoning often occurs in concert with other medical emergencies, such as smoke inhalation, and may affect many people at the same time.²⁹

Table 1 shows the variety of acute symptoms reported by patients after exposure to carbon monoxide in a number of clinical series.^{11,30,31} Patients often present with tachycardia and tachypnea, which are compensatory mechanisms for cellular hypoxia. Headache, nausea, and vomiting are common symptoms. Presyncope, syncope, and seizures may result from cellular hypoxia and cerebral vasodilatation, which can also lead to cerebral edema. Angina, pulmonary edema, and arrhythmias may result from increased cardiac output caused by cellular hypoxia, carbon monoxide-myoglobin binding, and diminished oxygen release.³²⁻³⁴ In patients with underlying pulmonary or cardiac disease, the symptoms of their disease may be worsened by impaired oxygen release.³⁵ The classic findings of cherry-red lips, cyanosis, and retinal hemorrhages occur rarely.³⁶ Erythematous lesions with bullae over bony prominences have been described but are not specific for carbon monoxide poisoning. Necrosis of the sweat glands is a characteristic histologic feature.³⁷

The severity of symptoms ranges from mild (constitutional symptoms) to severe (coma, respiratory depression, and hypotension). It is important to recognize that carboxyhemoglobin levels do not corre-

TABLE 1. ACUTE SYMPTOMS REPORTED BY 196 PATIENTS AFTER EXPOSURE TO CARBON MONOXIDE.*

SYMPTOM	PERCENTAGE OF PATIENTS
Headache	91
Dizziness	77
Weakness	53
Nausea	47
Difficulty in concentrating or confusion	43
Shortness of breath	40
Visual changes	25
Chest pain	9
Loss of consciousness	6
Abdominal pain	5
Muscle cramping	5

*Data are from Ely et al.,¹¹ Myers et al.,³⁰ and Burney et al.³¹

late well with the severity of symptoms in a substantial number of cases. The duration of exposure appears to be an important factor mediating toxicity. Being in a carbon monoxide-containing environment for one hour or more may increase morbidity.³⁸ If no dissolved carbon monoxide is present in the plasma, the symptoms can be minimal even with extremely high levels of carboxyhemoglobin, as experiments in animals show.²⁰ Therefore, the decision whether to administer hyperbaric oxygen therapy cannot be made only on the basis of carboxyhemoglobin levels.^{30,39}

DELAYED NEUROPSYCHIATRIC SYNDROME

Many patients with carbon monoxide poisoning do not have acute signs of cerebral impairment. Delayed onset of neuropsychiatric symptoms after apparent recovery from the acute intoxication has been described 3 to 240 days after exposure. The syndrome is estimated to occur in 10 to 30 percent of victims, but the reported incidence varies widely.⁴⁰⁻⁴² Symptoms such as cognitive and personality changes, parkinsonism, incontinence, dementia, and psychosis have been described.^{42,43} No clinical or laboratory results predict which patients are at risk for this complication, but advanced age appears to be a risk factor. Recovery from delayed neuropsychiatric syndrome occurs in 50 to 75 percent of affected persons within one year.⁴² Different abnormalities have been shown by computed tomography, molecular resonance imaging, and single-photon-emission computed tomography. The regions most commonly involved include the globus pallidus and the deep white matter.^{42,44}

Delayed neuropsychiatric sequelae after exposure

to carbon monoxide have been the subject of several reports.^{42,45} The mechanisms are uncertain, but hypoxia alone is not sufficient to explain the observed clinical manifestations. Postischemic reperfusion injury as well as the effects of carbon monoxide on vascular endothelium and oxygen-radical-mediated brain lipid peroxidation may also have a role.⁴⁶ In addition, nitric oxide liberated from platelets at the time of carbon monoxide exposure has been linked to central nervous system damage.⁴⁷

DIAGNOSIS

Because carbon monoxide poisoning has no pathognomonic signs or symptoms, a high level of suspicion, particularly among primary care clinicians and emergency medicine specialists, is essential for making the diagnosis. The measurement of carbon monoxide levels alone may be insufficient to rule out the diagnosis, but in the majority of cases, increased levels of carboxyhemoglobin will be diagnostic. Serum levels of carboxyhemoglobin may already have fallen substantially at the time of presentation to the emergency department. Therefore, elevated carbon monoxide values in the exhaled air of the patients or in the ambient air at the scene of exposure can help confirm the diagnosis. This latter test can be performed by fire departments and should be encouraged. Blood obtained on the scene by emergency medical technicians may also be helpful for confirming the diagnosis.

Venous blood samples are adequate for measurements of carboxyhemoglobin,⁴⁸ although arterial samples allow for the additional determination of coexisting acidosis. Carboxyhemoglobin has to be measured directly with a spectrophotometer. Pulse oximetry cannot distinguish carboxyhemoglobin from oxyhemoglobin at the wavelengths that are commonly employed by most oximeters (pulse-oximetry gap).^{49,50}

When the diagnosis of carbon monoxide poisoning has been established, a detailed neurologic examination and neuropsychological testing should be performed to document neurologic and neuropsychiatric abnormalities, which may be subtle. The Carbon Monoxide Neuropsychological Screening Battery is a frequently used tool that takes 30 minutes to administer and provides a base line for assessing subsequent changes in mental status.⁵¹ Computed tomographic imaging of the head is not helpful in establishing the diagnosis of carbon monoxide intoxication, but it may be used to rule out other conditions that might result in changes in mental status or loss of consciousness in patients presenting to an acute care facility.

TREATMENT

The carbon monoxide-intoxicated patient must first be removed from the source of carbon monoxide production without endangering the health of the

rescuing personnel. Firefighters must use breathing apparatus not only to supply oxygen but also to protect against carbon monoxide poisoning. High-flow oxygen, preferably 100 percent as normobaric oxygen, should be administered to the patient immediately. Oxygen shortens the half-life of carboxyhemoglobin by competing at the binding sites of hemoglobin and improves tissue oxygenation.⁵² Oxygen should be administered until the carboxyhemoglobin level has become normal. In patients with carbon monoxide poisoning who have been rescued from a fire, special consideration should be given to the respiratory status and the airway, since urgent or prophylactic intubation may be necessary.

Most patients can be evaluated and treated in an ambulatory setting. Hospitalization should be considered for patients with severe poisoning, serious underlying medical problems, or accompanying injuries. Patients often have concomitant problems, including smoke inhalation and burns, that require specialized treatment and may necessitate transfer to specialized facilities.

Since carbon monoxide may affect others who have been exposed to the same source, appropriate local agencies, usually the fire department, should be alerted to investigate the source of the intoxication and arrange for all other possible victims to be screened.

NORMOBARIC VERSUS HYPERBARIC OXYGEN

Carbon monoxide elimination is related to minute ventilation, the duration of exposure, and the fraction of inspired oxygen (FiO_2). The half-life of carboxyhemoglobin is 4 to 6 hours when the patient is breathing room air, 40 to 80 minutes when the patient is breathing 100 percent oxygen, and only 15 to 30 minutes when the patient is breathing hyperbaric oxygen.⁵³ In 1895 Haldane showed that hyperbaric oxygen prevented carbon monoxide poisoning in mice,⁵⁴ and since 1962 hyperbaric oxygen has been used to treat carbon monoxide poisoning.⁵⁵

The indications for hyperbaric-oxygen therapy have recently been reviewed in the *Journal*.⁵⁶ Hyperbaric-oxygen therapy hastens the resolution of symptoms. It is unclear whether hyperbaric-oxygen therapy influences the rate of late sequelae or mortality in non-life-threatening carbon monoxide poisoning, since different studies have led to conflicting conclusions.^{57,58} Coma is an undisputed indication for hyperbaric-oxygen therapy. Outcome studies of hyperbaric-oxygen therapy have not yet identified other circumstances in which this therapy is clearly indicated.⁵⁹ The indications for this therapy in patients with mild-to-moderate cerebral dysfunction are particularly disputed. Nonetheless, suggestions are available to help physicians decide whether to administer hyperbaric-oxygen therapy (Table 2).⁶⁰

TABLE 2. SUGGESTED INDICATIONS FOR HYPERBARIC-OXYGEN THERAPY IN PATIENTS WITH CARBON MONOXIDE POISONING.*

Coma
Any period of unconsciousness
Any abnormal score on the Carbon Monoxide Neuropsychological Screening Battery
Carboxyhemoglobin level >40%
Pregnancy and carboxyhemoglobin level >15%
Signs of cardiac ischemia or arrhythmia
History of ischemic heart disease and carboxyhemoglobin level >20%
Recurrent symptoms for up to 3 wk
Symptoms that do not resolve with normobaric oxygen after 4–6 hr

*Data are from Myers and Thom.⁶⁰

Once the diagnosis of carbon monoxide poisoning has been established, the physician must decide whether hyperbaric-oxygen therapy is indicated, and if so, make appropriate arrangements for a safe transfer to the nearest facility. More than 340 single-occupant chambers are available in the United States.⁶¹ Information on the location and use of decompression chambers is available by telephone from the Divers Alert Network at Duke University at 919-684-8111. Callers should request the Divers Alert Network on-call staff.

PREVENTION

Awareness of the dangers of carbon monoxide and public education are the keys to decreasing morbidity and mortality from carbon monoxide poisoning. Primary prevention is aimed at decreasing production of and exposure to carbon monoxide. The Environmental Protection Agency and the Occupational Safety and Health Administration provide regulations and suggestions,⁶²⁻⁶⁴ and general information is easily available from sources such as the American Gas Association. In particular, the current regulations of the Occupational Safety and Health Administration prohibit the exposure of workers to carbon monoxide levels exceeding 35 ppm, averaged over an 8-hour workday, with an upper limit of 200 ppm over a 15-minute period.⁶² Fuel-burning heating systems require regular professional maintenance and appropriate ventilation. Motor vehicles should not remain in enclosed spaces with the engine running, and the exhaust pipe must be free of obstructions (particularly snow and leaves). Outdoor gas grills should not be operated indoors. Media campaigns should warn the public about the dangers of carbon monoxide at times of increased risk, such as anticipated cold spells and snowstorms. Members of minority groups and non-English-speakers are at greatest

risk, and public education must be tailored to reach these parts of the population.^{65,66}

Secondary prevention efforts should be aimed at warning people about potentially harmful carbon monoxide concentrations in the environment. Although carbon monoxide detectors are inexpensive and widely available, they should not be considered a substitute for proper maintenance of appliances. There are currently no standard recommendations regarding their use in the home or the workplace.

We are indebted to Dr. Henry Koziel for his critical review of the manuscript.

REFERENCES

- Cobb N, Ertzel RA. Unintentional carbon monoxide-related deaths in the United States, 1979 through 1988. *JAMA* 1991;266:659-63.
- National Center for Health Statistics. Vital statistics of the United States, 1988. Washington, D.C.: Government Printing Office, 1991. (DHHS publication no. (PHS) 89-1102.)
- Geehr EC, Salluzzo R, Bosco S, Braaten J, Wahl T, Wallenkamp V. Emergency health impact of a severe storm. *Am J Emerg Med* 1989;7:598-604.
- Deaths from motor-vehicle-related unintentional carbon monoxide poisoning — Colorado, 1996, New Mexico, 1980–1995, and United States, 1979–1992. *MMWR Morb Mortal Wkly Rep* 1996;45:1029-32.
- Carbon monoxide poisonings associated with snow-obstructed vehicle exhaust systems — Philadelphia and New York City, January 1996. *MMWR Morb Mortal Wkly Rep* 1996;45:1-3.
- Carbon monoxide poisoning at an indoor ice arena and bingo hall — Seattle, 1996. *MMWR Morb Mortal Wkly Rep* 1996;45:265-7.
- Farrow JR, Davis GJ, Roy TM, McCloud LC, Nichols GR II. Fetal death due to nonlethal maternal carbon monoxide poisoning. *J Forensic Sci* 1990;35:1448-52.
- Hausberg M, Somers VK. Neural circulatory responses to carbon monoxide in healthy humans. *Hypertension* 1997;29:1114-8.
- Hee J, Callais F, Momas I, et al. Smokers' behaviour and exposure according to cigarette yield and smoking experience. *Pharmacol Biochem Behav* 1995;52:195-203.
- Meredith T, Vale A. Carbon monoxide poisoning. *BMJ* 1988;296:77-9.
- Ely EW, Moorhead B, Haponik EF. Warehouse workers' headache: emergency evaluation and management of 30 patients with carbon monoxide poisoning. *Am J Med* 1995;98:145-55.
- Baker SP, O'Neill B, Ginsburg MJ, Li G. The injury fact book. 2nd ed. New York: Oxford University Press, 1992.
- Stewart RD. The effect of carbon monoxide on humans. *Annu Rev Pharmacol* 1975;15:409-23.
- Stewart RD, Hake CL. Paint-remover hazard. *JAMA* 1976;235:398-401.
- Forbes WH, Sargent E, Roughton FJW. Rate of carbon monoxide uptake by normal men. *Am J Physiol* 1945;143:594-608.
- Coburn RF. The carbon monoxide body stores. *Ann N Y Acad Sci* 1970;174:11-22.
- Coburn RF, Forman HJ. Carbon monoxide toxicity. In: Farhi LE, Tenney SM, eds. *Handbook of physiology*. Section 3. The respiratory system. Vol. 4. Gas exchange. Bethesda, Md.: American Physiological Society, 1987:439-56.
- Rodkey FL, O'Neal JD, Collison HA, Uddin DE. Relative affinity of hemoglobin S and hemoglobin A for carbon monoxide and oxygen. *Clin Chem* 1974;20:83-4.
- Roughton FJW, Darling RC. The effect of carbon monoxide on oxy-hemoglobin dissociation curve. *Am J Physiol* 1944;141:17-31.
- Goldbaum LR, Ramirez RG, Absalon KB. What is the mechanism of carbon monoxide toxicity? *Aviat Space Environ Med* 1975;46:1289-91.
- Zhang J, Piantadosi CA. Mitochondrial oxidative stress after carbon monoxide hypoxia in the rat brain. *J Clin Invest* 1992;90:1193-9.
- Thom SR. Carbon monoxide-mediated brain lipid peroxidation in the rat. *J Appl Physiol* 1990;68:997-1003.
- Idem*. Dehydrogenase conversion to oxidase and lipid peroxidation in brain after carbon monoxide poisoning. *J Appl Physiol* 1992;73:1584-9.
- Longo LD, Hill EP. Carbon monoxide uptake and elimination in fetal and maternal sheep. *Am J Physiol* 1977;232:H324-H330.
- Norman CA, Halton DM. Is carbon monoxide a workplace teratogen? A review and evaluation of the literature. *Ann Occup Hyg* 1990;34:335-47.
- Ginsberg MD, Myers RE. Fetal brain injury after maternal carbon monoxide intoxication: clinical and neuropathologic aspects. *Neurology* 1976;26:15-23.
- Robkin MA. Carbon monoxide and the embryo. *Int J Dev Biol* 1997;41:283-9.
- Kales SN. Carbon monoxide intoxication. *Am Fam Physician* 1993;48:1100-4.
- Thom SR. Smoke inhalation. *Emerg Med Clin North Am* 1989;7:371-87.
- Myers RAM, Snyder SK, Emhoff TA. Subacute sequelae of carbon monoxide poisoning. *Ann Emerg Med* 1985;14:1163-7.
- Burney RE, Wu SC, Nemiroff MJ. Mass carbon monoxide poisoning: clinical effects and results of treatment in 184 victims. *Ann Emerg Med* 1982;11:394-9.
- Turino G. Effect of carbon monoxide on the cardiorespiratory system: carbon monoxide toxicity, physiology and biochemistry. *Circulation* 1981;63:253A-259A.
- Becker LC, Haak ED Jr. Augmentation of myocardial ischemia by low level carbon monoxide exposure in dogs. *Arch Environ Health* 1979;34:274-9.
- DeBias DA, Banerjee CM, Birkhead NC, Greene CH, Scott SD, Harter WV. Effects of carbon monoxide inhalation on ventricular fibrillation. *Arch Environ Health* 1976;31:42-6.
- Williams J, Lewis RW II, Kealey GP. Carbon monoxide poisoning and myocardial ischemia in patients with burns. *J Burn Care Rehabil* 1992;13:210-3.
- Hardy KR, Thom SR. Pathophysiology and treatment of carbon monoxide poisoning. *J Toxicol Clin Toxicol* 1994;32:613-29.
- Torne R, Soyler HP, Leb G, Kerl H. Skin lesions in carbon monoxide intoxication. *Dermatologica* 1991;183:212-5.
- Bogusz M, Cholewa L, Pach J, Mlodkowska K. A comparison of two types of acute carbon monoxide poisoning. *Arch Toxicol* 1975;33:141-9.
- Norkool DM, Kirkpatrick JN. Treatment of acute carbon monoxide poisoning with hyperbaric oxygen: a review of 115 cases. *Ann Emerg Med* 1985;14:1168-71.
- Hart IK, Kennedy PGE, Adams JH, Cunningham NE. Neurological manifestation of carbon monoxide poisoning. *Postgrad Med J* 1988;64:213-6.
- Sawa GM, Watson CPM, Terbrugge K, Chiu M. Delayed encephalopathy following carbon monoxide intoxication. *Can J Neurol Sci* 1981;8:777-9.
- Choi IS. Delayed neurologic sequelae in carbon monoxide intoxication. *Arch Neurol* 1983;40:433-5.
- Min SK. A brain syndrome associated with delayed neuropsychiatric sequelae following acute carbon monoxide intoxication. *Acta Psychiatr Scand* 1986;73:80-6.
- Zagami AS, Lethlean AK, Mellick R. Delayed neurological deterioration following carbon monoxide poisoning: MRI findings. *J Neurol* 1993;240:113-6.
- Seeger D, Welch L. Carbon monoxide controversies: neuropsychologic testing, mechanism of toxicity, and hyperbaric oxygen. *Ann Emerg Med* 1994;24:242-8.
- Thom SR. Leukocytes in carbon monoxide-mediated brain oxidative injury. *Toxicol Appl Pharmacol* 1993;123:234-47.
- Ischiropoulos H, Beers ME, Ohnishi ST, Fisher D, Garner SE, Thom SR. Nitric oxide production and perivascular nitration in brain after carbon monoxide poisoning in the rat. *J Clin Invest* 1996;97:2260-7.
- Touger M, Gallagher EJ, Tyrell J. Relationship between venous and arterial carboxyhemoglobin levels in patients with suspected carbon monoxide poisoning. *Ann Emerg Med* 1995;25:481-3.
- Vegförs M, Lennmarken C. Carboxyhemoglobinaemia and pulse oximetry. *Br J Anaesth* 1991;66:625-6.
- Buckley RG, Aks SE, Eshom JL, Rydman R, Schaidler J, Shayne P. The pulse oximetry gap in carbon monoxide intoxication. *Ann Emerg Med* 1994;24:252-5.
- Messiers LD, Myers RAM. A neuropsychological screening battery for emergency assessment of carbon-monoxide-poisoned patients. *J Clin Psychol* 1991;47:675-84.
- Iiano AL, Raffin TA. Management of carbon monoxide poisoning. *Chest* 1990;97:165-9.
- Pace N, Strajman E, Walker EL. Acceleration of carbon monoxide elimination in man by high pressure oxygen. *Science* 1950;111:652-4.
- Haldane J. The relation of the action of carbonic oxide to oxygen tension. *J Physiol* 1895;18:201-17.
- Smith G. The treatment of carbon monoxide poisoning with oxygen at two atmospheres absolute. *Ann Occup Hyg* 1962;5:259-63.
- Tibbles PM, Edelsberg JS. Hyperbaric-oxygen therapy. *N Engl J Med* 1996;334:1642-8.
- Raphael JC, Elkharrat D, Jars-Guincestre MC, et al. Trial of normobaric and hyperbaric oxygen for acute carbon monoxide intoxication. *Lancet* 1989;2:414-9.

- 58.** Thom SR, Taber RL, Mendiguren II, Clark JM, Hardy KR, Fisher AB. Delayed neuropsychologic sequelae after carbon monoxide poisoning: prevention by treatment with hyperbaric oxygen. *Ann Emerg Med* 1995;25:474-80.
- 59.** Tibbles PM, Perrotta PL. Treatment of carbon monoxide poisoning: a critical review of human outcome studies comparing normobaric oxygen with hyperbaric oxygen. *Ann Emerg Med* 1994;24:269-76.
- 60.** Myers RAM, Thom SR. Carbon monoxide and cyanide poisoning. In: Kindwall EP, ed. *Hyperbaric medicine practice*. Flagstaff, Ariz.: Best Publishing, 1994:357.
- 61.** *Directory of hyperbaric chambers, United States and Canada*. Bethesda, Md.: Undersea and Hyperbaric Medical Society, 1996.
- 62.** OSHA fact sheet: carbon monoxide poisoning. Washington, D.C.: Department of Labor, 1996.
- 63.** Fact sheet: revision of carbon monoxide standard. Ann Arbor, Mich.: National Vehicle and Fuel Emissions Laboratory, 1996. (Order no. EPA420-F-96005.)
- 64.** Air quality criteria for carbon monoxide. Cincinnati: National Center for Environmental Publications and Information, 1990. (Order no. EPA 600/890045-E.)
- 65.** Houck PM, Hampson NB. Epidemic carbon monoxide poisoning following a winter storm. *J Emerg Med* 1997;15:469-73.
- 66.** Wrenn K, Conners GP. Carbon monoxide poisoning during ice storms: a tale of two cities. *J Emerg Med* 1997;15:465-7.

Northern Arizona

SCOTT VALENT, M.D.