



Figure 1. Study Schematic and Dosing Schedule



- a. Subjects randomized to Treatment Arm A will receive R-CHOP in combination with epcoritamab for 6 cycles followed by 2 additional cycles of epcoritamab
- b. Subjects randomized to Treatment Arm B will receive R-CHOP for 6 cycles followed by 2 additional cycles of rituximab
- c. Subjects who complete treatment or prematurely discontinue the assigned treatment arm for reasons other than disease progression will proceed to Post-Treatment Follow-Up until 5 years after the first subject first dose
- d. Subjects who display disease progression or refuse Post-Treatment Follow-Up Visits will proceed to Survival Follow-Up



A Arm A

- ❖ R-CHOP administered for 6 cycles beginning on Day 1 of each cycle
 - Cycle 1-6 (once every 3 weeks) 21-day cycles**
 - Rituximab 375 mg/m² IV on Day 1
 - Cyclophosphamide 750 mg/m² IV on Day 1
 - Doxorubicin 50 mg/m² IV on Day 1
 - Vincristine 1.4 mg/m² IV on Day 1
 - Prednisone 100 mg PO on Days 1-5
- ❖ Epcoritamab administered for 6 cycles in step-up dosing as follows:
 - Cycle 1**
 - C1D1 (Week 1): 0.16 mg
 - C1D8 (Week 2): 0.8 mg
 - C1D15 (Week 3 - 1st full dose): 48 mg
 - Cycles 2-4 (once weekly) 21-day cycle**
 - C2D1 (Week 4): 48 mg
 - C2D8 (Week 5): 48 mg
 - C2D15 (Week 6): 48 mg
 - C3D1 (Week 7): 48 mg
 - C3D8 (Week 8): 48 mg
 - C3D15 (Week 9): 48 mg
 - C4D1 (Week 10): 48 mg
 - C4D8 (Week 11): 48 mg
 - C4D15 (Week 12): 48 mg
 - Cycles 5-6 (once every 3 weeks) 21-day cycles**
 - Day 1 of each cycle: 48 mg
- ❖ Epcoritamab (48mg) administered for 2 additional cycles beginning on Day 1 of cycle 7
 - Cycle 7-8 (once every 3 weeks) 21-day cycle**
 - Day 1 of each cycle: 48 mg

B Arm B

- ❖ R-CHOP administered for 6 cycles beginning on Day 1 of each cycle
 - Cycle 1-6 (once every 3 weeks) 21-day cycles**
 - Rituximab 375 mg/m² IV on Day 1
 - Cyclophosphamide 750 mg/m² IV on Day 1
 - Doxorubicin 50 mg/m² IV on Day 1
 - Vincristine 1.4 mg/m² IV on Day 1
 - Prednisone 100 mg PO on Days 1-5
- ❖ R (Rituximab) administered for 2 additional cycles beginning on Day 1 of cycle 7
 - Cycle 7-8 (once every 3 weeks) 21-day cycle**
 - Rituximab 375 mg/m² IV on Day 1

C = cycle; D = day; IV = intravenous(ly); PO = orally